

1. Granath C, Freiholtz D, Bredin F, Olsson C, Franco-Cereceda A, **Björck HM**. ASA is associated with a lower prevalence of ascending aortic aneurysm and a decreased aortic expression of COX2. *J Am Heart Assoc.* 2022; Apr 26; e024346. PMID: 35470674
<https://pubmed.ncbi.nlm.nih.gov/35470674/>
2. Vignac M, **Björck HM**, Olsson C, Eriksson MJ, Jouven X, Michos ED, Franco-Cereceda A, Eriksson P, Gaye B. Sex differences in aortopathy and valve diseases among patients undergoing cardiac surgery. *The Annals of Thoracic Surgery.* 2022; Mar 7:S0003-4975(22)00304-6. Online ahead of print. PMID: 35271843
<https://pubmed.ncbi.nlm.nih.gov/35271843/>
3. Vignac M, Ntika S, Olsson C, Franco-Cereceda, A, **Björck HM**. Metformin therapy is not associated with the lower prevalence of ascending aortic aneurysm in diabetic patients. *Eur J Cardiothorac Surg.* 2022; 24;61(2):388-392. PMID: 34676406
<https://pubmed.ncbi.nlm.nih.gov/34676406/>
4. Ntika S, Jois H, Lång K, Olsson C, Franco-Cereceda A, **Björck HM**, Krizhanovskii C. Elevated glucagon-like peptide-1 and a Th2 shift may support reduced prevalence of thoracic aortic aneurysm in patients with diabetes. *J cardiovasc dev dis.* 2021; 8(11): 143. PMID: 34821696.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8618274/>
5. Granath C, Mohamed SA, Olsson C, Grattan M, Mertens L, Franco-Cereceda A, **Björck HM**. Valve disease and aortopathy associations of bicuspid aortic valve phenotypes differ between men and women. *Open Heart.* 2021; 8:e001857. PMID: 34670833
<https://pubmed.ncbi.nlm.nih.gov/34670833/>
6. Ntika S, Tracy LM, Franco-Cereceda A, **Björck HM**, Krizhanovskii C. Syndecan-1 expression is increased in the aortic wall of patients with Type 2 Diabetes, but unrelated to elevated fasting plasma Glucagon like peptide-1. *Biomedicines* 2021; 20;9(6):697. PMID: 34203009
<https://pubmed.ncbi.nlm.nih.gov/34203009/>
7. Granath C, Noren H, **Björck H**, Simon N, Olesen K, Rodin S, Grinnemo KH, Österholm C. Characterization of Laminins in Healthy Human Aortic Valves and a Modified Decellularized Rat Scaffold. *Biores Open Access.* 2020; 7;9(1):269-278. PMID: 33376633
<https://pubmed.ncbi.nlm.nih.gov/33376633/>
8. Hussain S, Khan AW, Akhmedov A, Suades R, Costantino S, Paneni F, Caidahl K, Mohammed SA, Hage C, Gkolfos C, **Björck H**, Pernow J, Lund LH, Löscher TF, Cosentino F. Hyperglycemia Induces Myocardial Dysfunction via Epigenetic Regulation of JunD. *Circ Res.* 2020; 23;127(10):1261-1273. PMID: 32815777
<https://pubmed.ncbi.nlm.nih.gov/32815777/>
9. Maleki S, Cottrill KA, Poujade F-A, Bhattachariya A, Bergman O, Gådin JR, Simon N, Lundströmer K, Franco-Cereceda A, **Björck HM**, Chan SY, and Eriksson P. The mir-200 family regulates key pathogenic events in ascending aortas of individuals with bicuspid aortic valves. *J Intern Med.* 2019; 285(1):102-114. PMID: 30280445
<https://pubmed.ncbi.nlm.nih.gov/30280445/>
10. Gould RA*, Aziz H*, Woods CE*, Seman-Senderos MA, Sparks E, Preuss CL, Wünnemann F, Bedja D, Moats CR, McClymont SA, Rose R, Sobeira N, Ling H, MacCarrick G, Kumar AA, Luyck I, Cannaerts E, Verstraeten A, **Björck HM**, Lehsau A-C, Jaskula-Ranga V, Lauridsen H, Shah AA, Bennett CL, Ellinor P, Lin H, Isselbacher E, Lino Cardenas CL, Butcher J, Hughes GC, Lindsay ME, Baylor-Hopkins Center for Mendelian Genomics, MIBAVA Leducq Consortium, Mertens L Franco-Cereceda A, Verhagen J, Wessels M, Mohamed SA, Eriksson P, Mital S, Van

Laer L, Loeys BL, Andelfinger G, McCallion AS# Dietz HC#. ROBO4 Mutations Predispose Individuals to Bicuspid Aortic Valve and Thoracic Aortic Aneurysm. *Nat Gen.* 2019; 51(1):42-50. PMID: 30455415
<https://pubmed.ncbi.nlm.nih.gov/30455415/>

11. Ward-Caviness CK, Huffman JE, Evertt K, Germain M, Van Dongen J, Hill WD, Jhun MA, Brody JA, Ghanbari M, Du L, Roetker NS, de Vries PS, Waldenberger M, Gieger C, Wolf P, Prokisch H, Koenig W, O'Donnell CJ, Levy D, Liu C, Truong V, Wells PS, Trégouët DA, Tang W, Morrison AC, Boerwinkle E, Wiggins KL, McKnight B, Guo X, Psaty BM, Sotoodehnia N, Boomsma DI, Willemsen G, Ligthart L, Deary IJ, Zhao W, Ware EB, Kardia SLR, Van Meurs JBJ, Uitterlinden AG, Franco OH, Eriksson P, Franco-Cereceda A, Pankow JS, Johnson AD, Gagnon F, Morange PE, de Geus EJC, Starr JM, Smith JA, Dehghan A, **Björck HM**, Smith NL, Peters A. DNA methylation age is associated with an altered hemostatic profile in a multi-ethnic meta-analysis. *Blood.* 2019; 11:134(2):218. PMID: 30042098
<https://pubmed.ncbi.nlm.nih.gov/30042098/>
 12. Kostina A, **Björck H**, Ignatieva E, Irtyuga O, Uspensky V, Semenova D, Maleki S, Tomilin A, Moiseeva O, Franco-Cereceda A, Gordeev M, Faggian G, Kostareva A, Eriksson P, Malashicheva A. Notch, BMP and WNT/β-catenin network is impaired in endothelial cells of the patients with thoracic aortic aneurysm. *Atheroscler Suppl.* 2018; 35:e6-e13. PMID: 30172576
<https://pubmed.ncbi.nlm.nih.gov/30172576/>
 13. **Björck HM**, Du L, Pulignani S, Paloschi V, Lundströmer K, Kostina AS, Österholm C, Malashicheva A, Kostareva A, Evangelista A, Teixidó-Tura G, Maleki S, Franco-Cereceda A, Eriksson P, and MIBAVA Leducq Consortium. Altered DNA methylation indicates an oscillatory flow mediated epithelial-to-mesenchymal transition signature in ascending aorta of patients with bicuspid aortic valve. *Sci Rep.* 2018; 9:8(1):2777. PMID: 29426841
<https://pubmed.ncbi.nlm.nih.gov/29426841/>
 14. Strawbridge RJ, Silveira A, Hoed MD, Gustafsson S, Luan J, Rybin D, Dupuis J, Li-Gao R, Kavousi M, Dehghan A, Haljas K, Lahti J, Gådin JR, Bäcklund A, de Faire U, Gertow K, Giral P, Goel A, Humphries SE, Kurl S, Langenberg C, Lannfelt LL, Lind L, Lindgren CCM, Mannarino E, Mook-Kanamori DO, Morris AP, de Mutsert R, Rauramaa R, Saliba-Gustafsson P, Sennblad B, Smit AJ, Syvänen AC, Tremoli E, Veglia F, Zethelius B, **Björck HM**, Eriksson JG, Hofman A, Franco OH, Watkins H, Jukema JW, Florez JC, Wareham NJ, Meigs JB, Ingelsson E, Baldassarre D, Hamsten A; IMPROVE study group. Identification of a novel proinsulin-associated SNP and demonstration that proinsulin is unlikely to be a causal factor in subclinical vascular remodelling using Mendelian randomisation. *Atherosclerosis.* 2017; 266:196-204. PMID: 29040868
<https://pubmed.ncbi.nlm.nih.gov/29040868/>
 15. Gillis E, Kumar AA, Luyckx I, Preuss C, Cannaerts E, van de Beek G, Wieschendorf B, Alaerts M, Bolar N, Vandeweyer G, Meester J, Wünnemann F, Gould RA, Zhurayev R, Zerbino D, Mohamed SA, Mital S, Mertens L, **Björck HM**, Franco-Cereceda A, McCallion AS, Van Laer L, Verhagen JMA, van de Laar IMBH, Wessels MW, Messas E, Goudot G, Nemcikova M, Krebssova A, Kempers M, Saleminck S, Duijnhouwer T, Jeunemaitre X, Albuison J, Eriksson P, Andelfinger G, Dietz HC, Verstraeten A, Loeys BL, and Mibava Leducq Consortium. Candidate Gene Resequencing in a Large Bicuspid Aortic Valve-Associated Thoracic Aortic Aneurysm Cohort: *SMAD6* as an Important Contributor. *Front Physiol.* 2017; 13:8:400. PMID: 28659821
<https://pubmed.ncbi.nlm.nih.gov/28659821/>
- Corrigendum: Candidate Gene Resequencing in a Large Bicuspid Aortic Valve-Associated Thoracic Aortic Aneurysm Cohort: *SMAD6* as an Important Contributor. *Front Physiol.* 2017; 25:8:730. PMID: 28974934
<https://pubmed.ncbi.nlm.nih.gov/28974934/>

16. Maleki S, Kjellqvist S, Paloschi V, Magné J, Branca RM, Du L, Hultenby K, Petrini J, Fuxe J; MIBAVA Leducq Consortium, Lehtiö J, Franco-Cereceda A, Eriksson P, and **Björck HM**. Mesenchymal state of intimal cells may explain higher propensity to ascending aortic aneurysm in bicuspid aortic valves. *Sci Rep.* 2016; 25:6:35712. PMID: 27779199
<https://pubmed.ncbi.nlm.nih.gov/27779199/>
17. McLeod O, Silveira A, Gertow K, Gådin J, Bäcklund A, Sennblad B, Baldassarre D, Veglia F, Humphries SE, Tremoli E, de Faire U, **Björck HM**, Hamsten A, Öhrvik J, Strawbridge RJ on behalf of the IMPROVE study group. Genetic loci on chromosome 5 are associated with circulating levels of interleukin-5 and eosinophil count in a European population with high risk for cardiovascular disease. *Cytokine.* 2016; 81:1-9. PMID: 26821299
<https://pubmed.ncbi.nlm.nih.gov/26821299/>
18. Petrini J, Yousry M, Eriksson P, **Björck HM**, Rickenlund A, Franco-Cereceda A, Caidahl K, Eriksson MJ. Intima-media thickness of the descending aorta in patients with bicuspid aortic valve. *Int J Cardiol Heart Vasc.* 2016; 2:11:74-79 PMID: 28616529
<https://pubmed.ncbi.nlm.nih.gov/28616529/>
19. Paloschi V, Gådin JR, Khan S, **Björck HM**, Du L, Maleki S, Roy J, Lindeman J HM, Mohamed SA, Tsuda T, Franco-Cereceda A, and Eriksson P. Aneurysm development in patients with a bicuspid aortic valve is not associated with transforming growth factor- β activation. *Arterioscler Thromb Vasc Biol.* 2015; 35(4):973-80. PMID 25745062
<https://pubmed.ncbi.nlm.nih.gov/25745062/>
20. Forsell C, **Björck HM**, Eriksson P, Franco-Cereceda A and Gasser TC. Biomechanical properties of the thoracic aneurysmal wall; Differences between Bicuspid Aortic Valve (BAV) and Tricuspid Aortic Valve (TAV) patients. *Ann Thorac Surg.* 2014; 98(1):65-71. PMID: 24881863
<https://pubmed.ncbi.nlm.nih.gov/24881863/>
21. Maleki S[§], **Björck HM**[§], Folkersen L, Nilsson R, Renner J, Caidahl K, Franco-Cereceda A, Länne T, Eriksson P. Identification of a novel flow-mediated gene expression signature in patients with bicuspid aortic valve. *J Mol Med (Berl).* 2013; 91(1):129-39. [§]Equal contribution. PMID: 22903503
<https://pubmed.ncbi.nlm.nih.gov/22903503/>
22. **Björck HM**, Renner R, Maleki S, Nilsson SFE, Kihlberg J, Folkersen L, Karlsson M, Ebbers T, Eriksson P, Länne T. Characterization of shear-sensitive genes in the normal rat aorta identifies Hand2 as a major flow-responsive transcription factor. *PLoS ONE.* 2012;7(12):e52227. PMID 23284944
<https://pubmed.ncbi.nlm.nih.gov/23284944/>
23. Ljungberg LU, De Basso R, Alehagen U, **Björck HM**, Persson K, Dahlström U, Länne T. Impaired abdominal aortic wall integrity in elderly men carrying the angiotensin-converting enzyme D allele. *Eur J Vasc Endovasc Surg.* 2011; 42(3):309-16. PMID: 21570325
<https://pubmed.ncbi.nlm.nih.gov/21570325/>
24. **Björck HM**, Eriksson P, Alehagen U, De Basso R, Ljungberg LU, Persson K, Dahlström U, Länne T. Gender-specific association of the plasminogen activator inhibitor-1 4G/5G polymorphism with central arterial blood pressure. *Am J Hypertens.* 2011; 24(7):802-8. PMID: 21490692
<https://pubmed.ncbi.nlm.nih.gov/21490692/>
25. Ljungberg L, Alehagen U, Länne T, **Björck H**, De Basso R, Dahlström U, Persson K. The association between circulating angiotensin-converting enzyme and cardiovascular risk in the elderly: a cross-sectional study. *J Renin Angiotensin Aldosterone Syst.* 2011; 12(3):281-9. PMID: 21273224

<https://pubmed.ncbi.nlm.nih.gov/21273224/>

26. **Björck HM**, Lanne T, Alehagen U, Persson K, Rundqvist L, Hamsten A, Dahlström U, Eriksson P. Association of genetic variation on chromosome 9p21.3 and arterial stiffness. *J Intern Med.* 2009; 265(3):373-81. PMID: 19019192
<https://pubmed.ncbi.nlm.nih.gov/19019192/>
27. Wågsäter D, Zhu C, **Björck HM**, Eriksson P. Effects of PDGF-C and PDGF-D on monocyte migration and MMP-2 and MMP-9 expression. *Atherosclerosis.* 2009; 202(2):415-23. PMID: 18573494
<https://pubmed.ncbi.nlm.nih.gov/18573494/>